Sent: Sunday, May 5, 2019 7:33 PM

To: Cathy Marcinkevage - NOAA Federal

Cc: J. Stuart

Subject: Re: salvage density data for steelhead

Do either of you know if the salvage density modeling for steelhead is based on wild steelhead salvage? Or hatchery + wild? Significant for our "pop level" context and the blurb from CWF just seems to describe modeling for winter-run...

On Fri, May 3, 2019 at 2:08 PM Cathy Marcinkevage - NOAA Federal < cathy.marcinkevage@noaa.gov wrote:

Without looking hard, does this help?

https://drive.google.com/open?id=1UGtWInMLfn9pUhPDYq7WV4FJd r-BoW5

On Fri, May 3, 2019 at 1:20 PM Barbara Byrne - NOAA Federal < <u>barbara.byrne@noaa.gov</u>> wrote: Is there somewhere in the ROC BA or CWF BA or BiOp that provides documentation on how these modeled salvage numbers are generated, with (for WR) or without (for all others) normalization?

--

Barb Byrne

Fish Biologist
NOAA Fisheries West Coast Region
U.S. Department of Commerce
Office: 916-930-5612
barbara.byrne@noaa.gov
California Central Valley Office
650 Capitol Mall, Suite 5-100
Sacramento, CA 95814



Find us online

www.westcoast.fisheries.noaa.gov



--

Barb Byrne
Fish Biologist
NOAA Fisheries West Coast Region

U.S. Department of Commerce
Office: 916-930-5612
barbara.byrne@noaa.gov
California Central Valley Office
650 Capitol Mall, Suite 5-100
Sacramento, CA 95814



Find us online

www.westcoast.fisheries.noaa.gov

