

---

**From:** Eric Danner - NOAA Federal <eric.danner@noaa.gov>  
**Sent:** Monday, March 18, 2019 3:06 PM  
**To:** Evan Sawyer - NOAA Federal  
**Cc:** Cathy Marcinkevage - NOAA Federal  
**Subject:** Re: Seasonal habitat capacity  
**Attachments:** PA-COS\_byWYT.png

Here is another one by WYT.

Eric

On Fri, Mar 15, 2019 at 9:43 AM Evan Sawyer - NOAA Federal <[evan.sawyer@noaa.gov](mailto:evan.sawyer@noaa.gov)> wrote:  
Hey Eric,

Thanks for the information yesterday. Can you provide a comparison of habitat capacity (from the WRLCM) between the PA and COS for the following "seasons":

Winter (Dec 1 - March 1)

Spring (March 1 - May 15 [maybe needs to be June 1? or May 1?])

Summer [temp mgmt.] (May 15 [or June 1] - Nov 1)

Fall (Oct. 1 - Dec 1) <-- I realize there's overlap with Fall.

These "seasons" generally correspond to the timing of Reclamation's operational decisions, so it may be a way to describe any differences in the operations (PA to COS). If there's no difference that's useful information too.

Thanks,

Evan

--

Evan Bing Sawyer,  
Natural Resource Management Specialist  
NOAA Fisheries West Coast Region  
U.S. Department of Commerce  
Office: (916) 930-3656  
[Evan.Sawyer@noaa.gov](mailto:Evan.Sawyer@noaa.gov)  
[www.westcoast.fisheries.noaa.gov](http://www.westcoast.fisheries.noaa.gov)



--

Eric Danner, Ph.D.  
Supervisory Research Ecologist  
Fisheries Ecology Division, Southwest Fisheries Science Center  
110 McAllister Way  
Santa Cruz, CA 95060  
831-420-3917  
<http://swfsc.noaa.gov/>