

Agree with Sarah, I think the redd dewatering information would be useful but I also agree that if it's based on down-scaled (up-scaled?) monthly data it's less useful.

All, especially Barb: Line 37-39 and 40-48. For lines 37-39 (Stan and Lower SJR floodplain inundation), the result is area inundated based on the average monthly flow. For Lines 40-48 (tribs and bypasses using the SIT relationships), it is area suitable as rearing habitat.

First, I told ICF to prioritize Lines 37-38 to give us Stan results (because there isn't a lot for the Stan in other methods). Any disagreements? **No.**

No

Next, are there strong feelings on priority of tribs vs bypasses or vice versa for Lines 42-48? Or some tribs vs others? If you really want your division, speak up! **I would think Upper and Middle Sacramento, but would assume all. I would like more understanding from someone who knows about why we would value some and not other sections.**

I think prioritizing other data limited tribs over the Sacramento is fine but I would then prioritize the bypasses as they were largely left out of the effects analysis in the BA (limited information in Appendix D).

Finally, what format would you want to see results? Average acres per month? And by water year type? Or exceedance plots for each month and WYT? **I agree with the format that Barb mentioned.**

I more agree with Brian in that monthly exceedance plots are good. If time is a limitation I would focus on "all year types" as I'm not entirely sure how useful the WYT will be given WYT is not a factor/criteria used in Reclamations proposed ops (it isn't is it?). I guess WYT could be useful for binning effects.

Evan and Sarah : SALMOD. The SALMOD model is applied to the Upper Sac, with that broken into five (I think) reaches and results generated for each reach. I don't know that we need all reaches. Could you look at the CWF SALMOD description and analysis and tell me which reaches we need, and if all of them, which we need first /most? I think reach 5 may be the most useful. I can point you to the locations in the CWF BiOp Wed morning. **It looks like it goes from Keswick to RBDD, and is broken into 14 segments for temperaure, and 8 for spawning? I would say Kewick to Balls Ferry (Seg 1-7 and 1-4).**

Yeah I didn't find the "5" segments but I think we'll need all of them because of fall-run redd distribution (SRKW) which extends down to Bend Bridge. If we're prioritizing them I would mostly agree with Sarah but continue downstream a bit to include battle creek and some spring-run(Seg 1-10 and 1-5).

All for now. Thanks all!

Cathy

On Thu, Feb 28, 2019 at 2:31 PM Cathy Marcinkevage - NOAA Federal <cathy.marcinkevage@noaa.gov> wrote:

All --

Please see the attached list of WUA and IFIM related methods. This is in response to an initial request from us to ICF. The green rows are more specific to our request, and the rest are similar/related by weren't asked for specifically.

Please respond by noon tomorrow with an indication of methods that you would like completed for your division.

Note that the last column is ICF's take on the method -- if you have more "on the ground" or update knowledge that differs, you should defer to your own expertise.

Let me know if you have any questions.

Thanks!
Cathy

----- Forwarded message -----

From: **Ellis, Gregg** <Gregg.Ellis@icf.com>

Date: Thu, Feb 28, 2019 at 1:50 PM

Subject: Available models.xlsx

To: Cathy Marcinkevage - NOAA Federal (cathy.marcinkevage@noaa.gov)

<cathy.marcinkevage@noaa.gov>

--

Evan Bing Sawyer,
Natural Resource Management Specialist
NOAA Fisheries West Coast Region
U.S. Department of Commerce
Office: (916) 930-3656
Evan.Sawyer@noaa.gov
www.westcoast.fisheries.noaa.gov

